

THE FOLLOWING MEDICATIONS SHOULD NOT BE TAKEN PRIOR TO THE 14C-UREA BREATH TEST

EXCLUSION PERIOD	TYPE OF MEDICATION	GENERIC#	TRADE NAME
	Antacids		ALU-TAB Andrews Tums Antacid DeWitts Antacid Powder ENO Gastrogel Gaviscon Mylanta Rennie Titralac
	H2 Antagonists	Cimetidine	Magicul Tagamet
		Ranitidine	Ausran Ranital Ranoxyl Ulcaid Ultac Zantac
		Famotidine	Ausfam Pamacid Pepzan Famohexal Pepcidine
		Nizatidine	Nizac Tazac Tacidine
For 7 Days Prior to Test	Proton Pump Inhibitors	Lansopazole	Lanzopran Zoton Zopral
		Rabeprazole	Parzol Prabez Razit Pariet Rabzole
		Omeprazole	Acimax Meprazol Omepral Ome pro - GA Pemzo Ozmep Probitor
		Esomeprazole	Nexium
		Pantoprazole	Gastenz Panthron Panto Pantoloc Pantomac Somac Suvacid Ozpan Pantofast Pantozol Salpraz Sozol Torzole
For 14 Days Prior to Test	Cytoprotectives	Sucralfate	Carafate Ulcyte
For 30 Days Prior to Test	Antibiotics	Multiple actives	Nexium Hp7 Vimovo Probitor HP7
		Amoxicillin (Penicillins)	Abbecillin Amoxil Augmentin Cilamox Fisamox Moxiclav Ranmoxy Yomax Alphamox Bgramin Clavulin Clamoxyl Clavycillin Curam Maxamox
		Ampicillin (Penicillins)	Ampicyn Ibimicyn Austrapen
		Azithromycin (Macrolides)	Azith Zedd Zithromax Zitrocin
		Clarithromycin (Macrolides)	Clarac Clarithro Klacid Clarihexal Kalixocin
		Erythromycin (Macrolides)	EES E-Mycin Eryc Erythrocin
		Roxithromycin (Macrolides)	Blaxsig Roxar Roximycin Rulide
		Tetracyclines:	Akamin Doryx Doxsig Doxy Frakas Tetracycline# Doxylin Doxyhexal Minocycline# Minomycin Doxycycline# Vibra-tabs
		Cephalexin	Keflex Cephatrust Ibilex Cilex Ialex Rancef
		Cefaclor (Cephalosporins)	Ceclor Karlor Keflor Ozcef
		Ciprofloxacin (Quinolones)	C-Flox Ciprol Proquin Cifran Ciproxin
		Other Anti-infectives:	Metronidazole Tinidazole Rifaximin Bismuth Flagyl Metronide Fasigyn Xifaxan Denol Metrogyl Simplotan

- # Apply to all generic brand of the active ingredient listed.
- Information compiled in this list is taken from MIMS MAY/JUNE 2013.
- The above is NOT the exhausted list especially antibiotics and anti-infective drugs, if you unsure of the classification of the medication, please consult your doctor or pharmacist.
- Note: All antibiotics except Vancomycin, Sulfa (Alprim, Bactrim, Resprim, Septrin, etc) and Amphotericin B should be withheld for at least 4 weeks for maximal accuracy.